



For more information, contact the Neuro-Oncology Clinical Trial Team at: neuro.oncology@jwci.org 310-829-8265

Clinical Trial Investigators

Achal S. Achrol, MD
Garni Barkhoudarian, MD
Daniel Kelly, MD
Santosh Kesari, MD, PhD,
Steven O'Day, MD
Marlon Garzo Saria, PhD, RN

Clinical Trial Team

Najee Boucher, CRA najee.boucher@providence.org 310-582-7460

> Jaya Gill, RN, BSN jaya.gill@providence.org 310-582-7437

Annie Heng, RN, BSN HengA@jwci.org 310-582-7457

Tiffany Juarez, PhD Tiffany Juarez@jwci.org 310-449-5225

Anand Moses, CRA MosesA@jwci.org 310-582-7097

Sponsor

SanBio, Inc.

Saint John's Health Center

PROVIDENCE Health & Services

CLINICAL TRIAL ANNOUNCEMENT

Study of Modified Stem Cells (SB623) in Patients with Chronic Motor Deficit from Traumatic Brain Injury

Official Title: A Double-Blind, Controlled Phase 2 Study of the Safety and Efficacy of Modified Stem Cells (SB623) in Patients with Chronic Motor Deficit from Traumatic Brain Injury (TBI)

This is a double-blind, sham-surgery controlled study of stereotactic, intracranial injection of SB623 cells in patients with fixed motor deficits from traumatic brain injury.

Two groups, Group 1 (2.5, 5, and 10 million SB623 cells combined) and Group 2 (sham placebo), will be included in this study in a 3:1 randomization scheme. Group 1 will be further randomized in a 1:1:1 ratio to receive either 2.5 million SB623 cells, 5 million SB623 cells, or 10 million SB623 cells. Group 2 will receive a sham surgery at a 3:1 randomization ratio. Randomization will be performed via an interactive web/voice response system (IXRS), stratified by GOS-E) score (recorded in the IXRS at the clinical site).

SB623 cells are bone-marrow-derived stromal cells that have been transiently transfected with the intercellular domain of the human Notch-1 gene.

Key Inclusion Criteria:

- Age 18-75 years
- Documented history of TBI, with correlated findings by MRI or CT
- At least 12 months post-TBI
- GOS-E score of 3-6 (i.e. moderate or severe disability)

Key Exclusion Criteria:

- History or presence of any other major neurological disease
- Any seizures in the prior 3 months
- Presence of craniectomy (without bone flap replacement) or other contraindication to stereotactic surgery